НОВАЯ ЭРА НАУКИ И ЗНАНИЙ
МЕЖДУНАРОДНЫЙ НАУЧНЫЙ ЖУРНАЛ
global-science-review.com
editor@global-science-review.com

CLINICAL COURSE OF ERYSIPELAS IN YOUNG CHILDREN

Qodirova Marjona Oybek qizi
Bekmuratova Sarvinoz Komiljon qizi
Faculty of Pediatrics No. 2, Tashkent State Medical University, Tashkent, Uzbekistan
Scientific Supervisor: Mirzayeva Gulhayo Abdurashidovna
Assistant, Department of Infectious Diseases, Pediatric Infectious Diseases, Phthisiology and Pulmonology, Tashkent State Medical University, Tashkent, Uzbekistan

Abstract: Erysipelas is an acute bacterial infection of the skin and superficial lymphatic vessels, characterized by localized erythema, swelling, tenderness, and clearly demarcated inflammatory borders. Although erysipelas is more frequently reported in adults, it may also occur in young children and requires particular clinical attention because of their age-related physiological characteristics and the potential for rapid progression of infectious diseases. This article examines the main clinical manifestations, pathogenesis, diagnostic approaches, treatment principles, possible complications, and preventive measures associated with erysipelas in young children. Particular attention is given to fever, systemic manifestations of infection, the localization and progression of skin lesions, and the importance of distinguishing erysipelas from other pediatric skin and soft-tissue infections. Early recognition, appropriate antimicrobial therapy, careful clinical monitoring, and timely identification of complications are essential for achieving favorable outcomes. An individualized approach based on the child's age, clinical severity, and general condition is recommended.

Keywords: erysipelas, young children, bacterial infection, skin and soft-tissue infections, streptococci, clinical manifestations, diagnosis, antimicrobial therapy, complications, prevention.

Introduction

Erysipelas is an acute infectious disease involving the superficial layers of the skin and the superficial lymphatic vessels. It is characterized by the sudden appearance of a localized inflammatory lesion, usually accompanied by redness, swelling, increased skin temperature, and tenderness. The disease is most commonly associated with beta-hemolytic streptococci, particularly group A Streptococcus (*Streptococcus pyogenes*). Less frequently, other streptococcal groups may be involved.

In pediatric practice, bacterial skin and soft-tissue infections represent an important clinical problem because their manifestations may develop rapidly and sometimes resemble other infectious or inflammatory conditions. Young children may be unable to describe pain, discomfort, or other symptoms accurately. Consequently, physicians must rely on careful physical examination, observation of behavioral changes, assessment of body temperature, and evaluation of the progression of skin lesions.

Erysipelas generally develops when pathogenic microorganisms enter the skin through a damaged area. Small wounds, scratches, insect bites, excoriations, and inflammatory skin conditions may provide an entry point for infection. In some cases, the initial skin defect is difficult to identify. After entering the tissues, bacteria trigger an inflammatory response involving the superficial dermis and lymphatic vessels.

The clinical presentation of erysipelas varies according to the anatomical location, extent of inflammation, age of the child, and severity of the infectious process. The face and lower extremities are commonly affected, although lesions may also develop in other anatomical regions. In infants, particular attention should be paid to inflammatory changes around the umbilicus and diaper area.

The importance of studying erysipelas in young children lies in the need for early diagnosis, appropriate treatment, and prevention of potentially serious

complications. Delayed recognition may allow the infection to progress and increase the risk of systemic illness. Therefore, understanding the characteristic clinical features and distinguishing erysipelas from other skin infections are essential components of pediatric clinical practice.

The aim of this article is to describe the clinical course of erysipelas in young children, identify its principal manifestations, discuss diagnostic approaches, and summarize current principles of treatment and prevention.

Materials and Methods

This article is based on a narrative review of scientific literature and established clinical guidance concerning erysipelas and bacterial skin and soft-tissue infections. Relevant information was considered from infectious disease guidelines, pediatric clinical reviews, and professional medical references addressing the etiology, clinical manifestations, diagnosis, treatment, and complications of erysipelas.

The review focused on the characteristic symptoms observed in young children, the mechanisms of infection, the identification of clinical severity, differential diagnosis, and principles of antimicrobial treatment. Particular emphasis was placed on the need to adapt clinical assessment and management to the child's age and general condition.

The article represents a literature-based review and does not report original patient recruitment, individual clinical observations, laboratory results, or statistical analysis. Accordingly, no patient cohort or original experimental findings are presented.

Results and Discussion

Etiology and Pathogenesis of Erysipelas

The principal causative agent of erysipelas is group A beta-hemolytic Streptococcus. These bacteria can enter the body through disruptions in the skin

barrier and initiate an acute inflammatory response. Other beta-hemolytic streptococci may occasionally cause similar infections.

The integrity of the skin is an important protective mechanism against bacterial invasion. In young children, this barrier may be compromised by minor injuries, scratching, eczema, diaper-associated irritation, insect bites, or other inflammatory skin conditions. Once microorganisms penetrate the skin, they may spread through the superficial dermis and lymphatic vessels, producing the characteristic clinical features of erysipelas.

The inflammatory response causes dilation of local blood vessels, increased vascular permeability, and accumulation of fluid in the affected tissues. These processes result in redness, edema, warmth, and tenderness. The release of inflammatory mediators may also contribute to fever, malaise, irritability, and other systemic manifestations.

The severity of the disease depends on several factors, including the extent of bacterial invasion, the child's immune response, the anatomical site involved, and the presence of underlying conditions. The clinical condition of infants and younger children requires particularly careful assessment because systemic symptoms may be nonspecific.

Erysipelas should be distinguished from cellulitis. Erysipelas typically affects more superficial tissues and produces a raised, sharply demarcated inflammatory border. Cellulitis generally involves deeper layers of the dermis and subcutaneous tissue, with less clearly defined margins. However, considerable clinical overlap exists, and the distinction may not always be straightforward.

Clinical Manifestations in Young Children

The onset of erysipelas is often acute. Fever, chills, irritability, reduced activity, and general discomfort may appear before or alongside the visible skin lesion. In young children, decreased appetite, disturbed sleep, excessive crying, and reduced interest in normal activities may be early indicators of illness.

The local manifestations are usually the most characteristic features. The affected area becomes red, warm, swollen, and tender. The inflammatory lesion often has clearly defined, slightly elevated borders, which distinguish it from many other inflammatory skin conditions. The intensity of redness may vary depending on the child's skin pigmentation and the stage of the disease.

The affected region may expand over a relatively short period. Parents may notice that the skin becomes increasingly swollen, warm, or painful. In more severe cases, blisters may develop over the inflamed area. Bullous lesions require careful evaluation because they may indicate a more complicated presentation or an alternative diagnosis.

Fever is an important systemic manifestation. Its intensity varies from mild elevation of body temperature to high fever accompanied by marked weakness and irritability. Tachycardia, poor feeding, vomiting, or reduced fluid intake may accompany more significant illness. These findings are not specific to erysipelas but can help determine the overall severity of the infection.

The location of the lesion influences the clinical presentation. Facial erysipelas may produce noticeable swelling and redness around the cheeks, nose, or other facial structures. Periorbital involvement is particularly important because infection in this region may be associated with deeper orbital disease, which requires urgent assessment. Lesions affecting the extremities may cause swelling and discomfort during movement.

In infants, erysipelas may occur around the umbilicus or in the diaper region, where irritation and minor skin damage can facilitate bacterial entry. Redness in these areas should not automatically be attributed to simple skin irritation, particularly when accompanied by fever, spreading inflammation, or changes in the infant's general condition.

The clinical course may be mild, moderate, or severe. Mild disease generally presents with localized inflammation and limited systemic symptoms. Moderate

disease may involve more extensive skin lesions, significant fever, and reduced activity. Severe presentations may be associated with rapidly progressive inflammation, marked systemic illness, circulatory instability, or other signs of invasive infection.

Clinical severity cannot be determined by the size of the skin lesion alone. The child's age, level of consciousness, hydration, respiratory status, circulation, and ability to feed must also be considered.

Diagnostic Approaches

The diagnosis of erysipelas is primarily clinical and is based on the patient's history and physical examination. The physician should determine the time of symptom onset, the rate of lesion progression, the presence of fever, possible skin injuries, previous episodes, and any recent antimicrobial treatment.

Physical examination should assess the location, size, color, warmth, tenderness, and borders of the lesion. The presence of edema, blisters, purulent discharge, skin discoloration, or necrotic changes should be documented. The clinician should also assess regional lymph nodes and look for possible portals of bacterial entry.

General clinical assessment is essential in young children. Body temperature, heart rate, respiratory rate, hydration status, level of activity, and feeding tolerance should be evaluated according to age-appropriate standards. Signs of systemic toxicity require urgent medical attention.

Laboratory investigations are selected according to the severity of the condition and the child's clinical presentation. A complete blood count may demonstrate leukocytosis with neutrophil predominance. C-reactive protein and other inflammatory markers may support the assessment of systemic inflammation and help monitor the clinical course. However, these findings are nonspecific and cannot independently establish the diagnosis.

Blood cultures are not routinely required in every uncomplicated case. They may be indicated in children with severe systemic illness, suspected bacteremia, immunocompromising conditions, or other circumstances in which identifying the causative organism would influence management. Microbiological samples from skin lesions are considered when there is purulent material, an unusual exposure, suspected resistant organisms, or an atypical clinical course.

Imaging is not routinely necessary for typical superficial erysipelas. Ultrasonography or other imaging methods may be appropriate when an abscess, deeper infection, or another complication is suspected.

Differential diagnosis is an important part of clinical assessment. Erysipelas may resemble cellulitis, impetigo, allergic contact dermatitis, insect-bite reactions, urticaria, and other inflammatory skin conditions. Facial swelling may also require differentiation from angioedema and deeper facial or orbital infections.

A particularly important diagnostic consideration is necrotizing soft-tissue infection. Rapid progression, severe pain disproportionate to the visible lesion, skin necrosis, bullae, altered sensation, or systemic deterioration should prompt immediate evaluation for a potentially life-threatening condition. In such circumstances, urgent specialist assessment must not be delayed by routine investigations.

Principles of Treatment

The primary treatment of erysipelas is appropriate antimicrobial therapy directed against the likely causative bacteria. Penicillin-based treatment is commonly used for uncomplicated streptococcal erysipelas, while alternative agents may be necessary in children with documented serious allergies or when the clinical presentation suggests a different pathogen.

The selection of an antimicrobial agent, route of administration, dosage, and duration must be determined by a qualified clinician according to the child's age, body weight, clinical severity, allergy history, local resistance patterns, and

applicable pediatric guidelines. Treatment should never be initiated or modified by caregivers without medical advice.

Oral antimicrobial therapy may be appropriate for selected children who are clinically stable, can tolerate oral medication, and have no indications for hospital admission. Parenteral treatment and hospital assessment may be required for severe illness, rapidly progressive lesions, systemic toxicity, inability to maintain adequate hydration, suspected complications, or other significant clinical concerns.

Supportive treatment is also important. Adequate fluid intake should be maintained, and age-appropriate antipyretic or analgesic medication may be used under medical guidance. The affected area should be protected from additional trauma. When an extremity is involved, elevation may help reduce swelling if tolerated by the child.

The affected skin should be observed regularly to assess whether the inflammatory border is expanding or the swelling is increasing. Marking the edge of the redness, when appropriate, can help healthcare professionals monitor progression. Caregivers should follow the treatment plan and seek reassessment if symptoms worsen or fail to improve as expected.

Clinical improvement is generally reflected by a reduction in fever, improved activity and appetite, and stabilization of the skin lesion. Local redness and swelling may resolve more slowly than systemic symptoms. Persistent or worsening inflammation should prompt reconsideration of the diagnosis, adherence to treatment, the possibility of an abscess, or a deeper infection.

Complications and Prognosis

Most cases of erysipelas respond favorably to timely and appropriate treatment. Nevertheless, complications may develop, particularly when infection progresses rapidly or systemic illness is present.

Local complications include blister formation, tissue damage, and, in uncommon severe cases, necrosis. Secondary abscess formation or extension into

deeper tissues may occur and should be considered when the lesion fails to respond as expected.

Systemic complications include bacteremia, sepsis, and, less commonly, infection involving deeper structures such as bone or joints. Facial infections require special attention because extension into adjacent anatomical structures may result in serious complications. Severe systemic symptoms, reduced consciousness, circulatory abnormalities, or rapidly worsening clinical status require immediate medical assessment.

The prognosis depends on early recognition, the appropriateness of treatment, the extent of infection, and the child's general health. In uncomplicated cases, fever and systemic symptoms often begin to improve after effective therapy, while complete resolution of skin changes may take longer.

Follow-up is important for confirming recovery and identifying persistent or recurrent infection. Repeated episodes should prompt evaluation for predisposing factors, including chronic skin inflammation, recurrent skin trauma, or other conditions that impair the skin barrier.

Prevention of Erysipelas in Young Children

Prevention is based primarily on maintaining skin integrity, recognizing early signs of infection, and providing appropriate care for minor skin injuries. Parents and caregivers should keep the child's skin clean, manage eczema and other inflammatory skin conditions according to medical advice, and prevent excessive scratching where possible.

Minor wounds should be cleaned appropriately and monitored for increasing redness, swelling, pain, or discharge. Diaper areas should be kept clean and dry, with measures taken to minimize persistent irritation. Umbilical skin in newborns should receive routine care according to established neonatal recommendations.

Good hand hygiene and appropriate infection-control practices are important in healthcare and childcare settings. Caregivers should avoid applying unprescribed antibiotics, irritating substances, or unverified traditional remedies to inflamed skin.

Parents should be informed about warning signs that require medical assessment, including fever associated with spreading redness, increasing tenderness, significant swelling, poor feeding, unusual sleepiness, and rapid deterioration. Infants with fever or suspected bacterial skin infection require prompt professional evaluation, with particular urgency in very young infants.

Education of parents and healthcare workers can improve early recognition and reduce delays in treatment. Preventive care should focus on early detection, appropriate management of skin injuries, and timely referral when the child's condition suggests a more serious infection.

Conclusion

Erysipelas in young children is an acute bacterial infection of the superficial dermis and lymphatic vessels that requires careful clinical assessment. Its characteristic manifestations include localized redness, warmth, swelling, tenderness, and sharply demarcated inflammatory borders. Fever, irritability, reduced appetite, and decreased activity may accompany the local skin changes.

Diagnosis is primarily based on clinical findings, supported by laboratory investigations when indicated. Differentiation from cellulitis, allergic skin reactions, abscesses, and potentially life-threatening deep infections is essential. Treatment depends on clinical severity and generally involves appropriate antimicrobial therapy, supportive care, and monitoring of the child's response.

Although the prognosis is favorable in most uncomplicated cases, delayed diagnosis may increase the risk of local and systemic complications. Early medical assessment, age-appropriate treatment, careful follow-up, skin care, and parent education are therefore central to effective management.

Further pediatric clinical research would be valuable for clarifying age-specific patterns of presentation, identifying factors associated with severe disease, and improving strategies for early diagnosis and prevention. A systematic approach to recognizing warning signs and initiating appropriate treatment can contribute to better outcomes in young children with erysipelas.

References

1. Stevens D. L., Bisno A. L., Chambers H. F., et al. Practice Guidelines for the Diagnosis and Management of Skin and Soft Tissue Infections: 2014 Update by the Infectious Diseases Society of America. *Clinical Infectious Diseases*. 2014;59(2):e10–e52. doi:10.1093/cid/ciu444.
2. Merck Manual Professional Edition. Erysipelas. Professional medical reference. Available at:https://www.merckmanuals.com/professional/infectious-diseases/bacterial-skin-infections/erysipelas
3. DermNet New Zealand. Erysipelas. Available at:https://dermnetnz.org/topics/erysipelas
4. Médecins Sans Frontières. MSF Medical Guidelines: Erysipelas and Cellulitis. Available at:https://medicalguidelines.msf.org/en/viewport/CG/english/erysipelas-and-cellulitis-16689672.html
5. Sartelli M., Guirao X., Hardcastle T. C., et al. 2018 WSES/SIS-E consensus conference: recommendations for the management of skin and soft-tissue infections. *World Journal of Emergency Surgery*. 2018;13:58. doi:10.1186/s13017-018-0219-9.